

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Rapid Fire 800 SG

HERBICIDE

ACTIVE CONSTITUENT: 800 g/kg GLYPHOSATE (present as the acid and ammonium salt)

GROUP M HERBICIDE

For the control of annual and perennial and aquatic weeds in many situations as per the Directions for Use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT



Indigo Specialty Products Pty Ltd
ABN 15 631 459 660
Unit 3 / 2 Ant Road
Yatala Qld 4207
Ph: 0402 725 887

Net Contents
10 kg



Rapid Fire 800 SG

HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Do not contaminate seed, feed or foodstuff. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

May irritate the nose and throat.

Corrosive. Attacks eyes and skin. Avoid contact with eyes and skin. If dust is present, wear disposable face mask covering mouth and nose. When opening the container, and preparing the spray, wear cotton overalls buttoned to the neck and wrist and elbow-length chemical resistant gloves and goggles. If applying by hand, wear cotton overalls buttoned to the neck and wrist and elbow-length chemical resistant gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce RAPID FIRE 800 SG Herbicide is available from Indigo Specialty Products on request.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use.

No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

	IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE
	SPECIALIST ADVICE IN AN EMERGENCY ONLY 0417 653 654 ALL HOURS – AUSTRALIA WIDE



10 kg

Batch No.

Date of
Manufacture

APVMA Approval No.: 68210/121825

8. GENERAL USES

Table with 4 columns: Situation, Weeds, Rate /ha, Critical Comments. Rows include Agricultural Areas, Forests, Pasture, and various crop applications like Cotton, Onions, Sugar Cane, and Cotton Pre-Harvest.

9. CONSERVATION TILLAGE

Table with 4 columns: Situation, Weeds, Rate /ha, Critical Comments. Rows include Southern Australia Prior to Sowing a Winter Crop or Pasture and Southern Australia Prior to Sowing a Winter Crop or Pasture.

Main table with 4 columns: Situation, Weeds, Rate /ha, Critical Comments. Rows include Northern Australia For Weed Control, POA Tussock Infested Pasture, Bent Grass Infested Pasture, Pasture Topping, Pasture Manipulation, Sorghum Control, Rice Direct Drilling, and Tasmania Only.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

This product is a non-volatile, water soluble granule with non-selective herbicidal activity against many annual and perennial broadleaf weeds and grasses.

Visible effects on annual weeds take 3-7 days, but on perennial weeds may not be obvious for 2-3 weeks or longer in some cases.

CROP ESTABLISHMENT

This product is recommended for control of emerged weeds prior to crop establishment. Suitable cultivation and/or sowing operations are required to provide seedbed conditions.

MIXING INSTRUCTIONS

This granular product dissolves readily in water. Where product is supplied in Water Soluble Measure Packs, these will readily break up in the spray tank.

For boom applications, water volumes should not be less than 6 litres per 1kg of RAPID FIRE 800 SG.

Ensure the spray tank is free of any residue of previous spray materials.

Fill the spray tank with one half the required amount of clean water and add the proper amount of this product.

TANK MIXTURES

This product may be tank-mixed with the following herbicides, insecticides and additives where recommended.

Mixing Instructions for All Tank Mixtures

- 1. Fill the spray tank 1/3 to 1/2 full with clean water and start agitation.
2. Add recommended herbicide/insecticide/additive to the spray tank and mix thoroughly.

TANK MIXTURES - HERBICIDES

Atrazine flowable* (DO NOT apply the tank-mix for Barnyard grass control), Farmalinx 2,4-D ester, Dicam, Tufman, Farmalinx Simazine, sulfometuron, pendimethalin, Metfluron 600 WG, Triflora, Farmalinx UVE MCPA 500 or 570, and Oxyfan 240.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use.

CAUTION

KEEP OUT OF REACH OF CHILDREN. READ SAFETY DIRECTIONS BEFORE OPENING OR USING. Do not spray near children, pets, or livestock.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Apply treatments using boom or Micronair equipment using a spray volume not less than 20 L/ha and using settings to produce a MEDIUM spray quality.

Product packaging for Indigo Specialty Products Rapid Fire 800 SG Herbicide. Includes 'CAUTION' warning, 'GROUP M HERBICIDE' label, and 'ACTIVE CONSTITUENT: 800 g/kg GLYPHOSATE (present as the acid and ammonium salt)'.

APPLICATION CHECKLIST

- Do not treat weeds under poor growing or dormant conditions (such as occur in drought, waterlogging, disease, insect damage or following frosts) as reduced weed control may result.
Do not add additional surfactants, or mix with any other agricultural chemicals, herbicide oils, or any other materials except as specifically directed on this label.

BOOM EQUIPMENT

Recommended rates of this product in 75 to 200 litres of clean water per hectare. When using this product at 250-700 g/ha spray volumes in the range 25-100 L/ha are preferred.

HIGH VOLUME APPLICATION (e.g. Knapsack and Handgun Equipment)

Adjust equipment to achieve an even spray pattern; for handgun equipment a D6 spray tip (Spraying Systems Australia P/L) or equivalent and an operating pressure of 400-700 kPa are recommended.

LOW VOLUME APPLICATION (e.g. Gas Gun or Splatter Gun)

Apply as even spray to cover all foliage. Refer to Weeds Controlled for the dilution and volume of mixture to be applied.

WIPER EQUIPMENT

Wiper equipment (e.g. ropewick, canvas, flat or carpet applicators) may be used to apply this product on to weeds growing in pasture and non-crop areas.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Apply contact with foliage, green stems or fruit of crops, desirable plants and trees, since severe injury or destruction may result.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS & ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. When controlling weeds near water, refer to label directions to minimise the entry of spray into the water.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Do not contaminate seed, feed or foodstuff.

SAFETY DIRECTIONS

May irritate the nose and throat. Corrosive. Attacks eyes and skin. Avoid contact with eyes and skin. If dust is present, wear disposable face mask covering mouth and nose.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce RAPID FIRE 800 SG Herbicide is available from Indigo Specialty Products on request.

Vertical text on the right edge of the page, including 'GROUP M HERBICIDE' and 'Indigo Specialty Products Pty Ltd'.