

Tombstone Disease Control

Fungicide USAGE CARD



- > A fungicide for use on Lawns, Azaleas, Roses, other general ornamental garden plants.
- > Provides systemic and contact disease control, both on a preventative and curative basis.
- > Controls Anthracnose, Brown Patch, Winter Fusarium, Leptosphaerulina (Leaf Blight), Curvularia (Fading Out) and Dollar Spot in lawn situations.
- > Controls azalea petal blight, myrtle rust, Botrytis Blight, Leaf Spots as well as mildew.

How ProForce Tombstone Disease Control Works

Tombstone Disease Control is a broad spectrum fungicide containing the active ingredients Trifloxystrobin (100g/L) and Tebuconazole (200g/L). One active ingredient is systemic in the plant and moves to the leaf tips once it enters plant parts. The other active ingredient bonds strongly onto plant surfaces providing long lasted, weather protected disease control.

Key Application Information

- > Avoid mowing lawns for 7 days after application.
- > Avoid spray application when rain is expected within 12 hours.
- > Avoid application when the soil is dry and plants are suffering from moisture stress.
- > Do not spray when shade temperatures are likely to exceed 30 °C.
- > Avoid application on edible plants in the garden.

Application Rates and Instructions

Lawns: Use 20mL – 30mL of Tombstone Disease Control in 10L of water per 100m². For preventive control, apply every 21-28 days when conditions favour disease development (e.g. warm, rainy weather). For curative control, apply when disease symptoms first appear and repeat every 14-21 days. For Brown patch control use the watering can method to ensure product is washed into the crown and root zone.

Azalea Petal Blight: Use 7.5mL of Tombstone Disease Control per 10 Litres of water. Apply thoroughly every two weeks from the expanded bud stage (before buds colour) until the end of flowering.

Black Spot and Powdery Mildew on Roses: Use 7.5mL – 15mL of Tombstone Disease Control per 10 Litres of water. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves. Apply as a preventative application 7-14 days apart when conditions favour disease development (e.g. warm, rainy weather), or as a curative treatment every 7-14 days after the onset of disease.

Botrytis Blight, Grey Mould, Downy Mildew and Leaf Spots on Ornamental Garden Plants: Use 7.5mL of Tombstone Disease Control per 10 Litres of water. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves. Apply as a curative treatment every 7-14 days after the onset of disease.

Myrtle Rust: Use 7.5mL – 15mL of Tombstone Disease Control per 10 Litres of water. Apply at first sign of disease or when conditions favour disease development. **Curative:** For existing infections, apply 2 applications at a 14-day interval. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves. **Preventative:** Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 14 days according to infection pressure.

Dollar Spot

