

Topdress Iron Xtra 21-2-5 (21-5-6) 4-5 Month

GUARANTEED ANALYSIS

40/11/11/12D /11/12/010		
TOTAL NITROGEN (N)**	21.00%	
3.00% Ammoniacal Nitrogen		
18.00% Urea Nitrogen**		
TOTAL PHOSPHORUS (P)	2.45%	
TOTAL POTASSIUM (K)	5.19%	
Magnesium (Mg)	0.50%	
Sulphur (S)	2.40%	
Copper (Cu)	0.06%	
Iron (Fe)	6.00%	
Manganese (Mn)	0.06%	
Molybdenum (Mo)	0.006%	
Zinc (Zn)	0.06%	

Derived From: Polymer-Coated Urea, Monoammonium Phosphate, Ammonium Sulphate, Sulphate of Potash, Muriate of Potash, Magnesium Carbonate, Magnesium Oxide, Magnesium Sulphate, Copper Oxide, Copper Sulphate, Ferric Oxide, Ferrous Sulphate, Iron Sucrate, Manganese Oxide, Manganese Sulphate, Sodium Molybdate, Zinc Oxide and Zinc Sulphate.

DESCRIPTION: UP TO 4 TO 5 MONTH ■ PROFESSIONAL CONTROLLED RELEASE FERTILISER FOR TOPDRESSING. BROADCAST, OR INCORPORATION APPLICATION OF TREES AND SHRUBS.

SOIL / MEDIA TEMPERATURE RELEASE RATES

 $10.0^{\circ}\text{C} / 50^{\circ}\text{F} = 6-7 \text{ month}$ $15.5^{\circ}C / 60^{\circ}F = 5-6 \text{ month}$ $21.0^{\circ}C / 70^{\circ}F = 4-5 \text{ month}$

 $26.5^{\circ}C / 80^{\circ}F = 3-4 \text{ month}$

BENEFITS

- 90% of the nitrogen is coated slow release for predictable
- Release of nitrogen is by diffusion.
- Release of nitrogen is not significantly affected by media type, moisture level, pH, or microbial activity.

APPLICATION RATES (Call for rates on larger containers)

Use LOW rate for low feeding, sensitive plants or under high soil temperatures. Use MEDIUM rate for medium to moderately heavy feeding plants.

Use HIGH rate only for heavy feeding hardy plants.

These application rates are based on the average temperature at the fertiliser location of 70°F (21.0°C).

Increase fertiliser application rates by 20% if average monthly temperatures are lower than 60°F (15.5°C). Lower application rates by 20% if average monthly temperatures are greater than 80°F (26.5°C).

TOPDRESS CONTAINER

Plant Nutrient Requirements / Uniformly apply (topdress) product onto the container surface using the amounts listed below.

DIAMETER (mm)	LOW	MEDIUM	HIGH
100 mm	0.5 g	1.0 g	1.5 g
125 mm	1.0 g	2.0 g	3.0 g
150 mm	2.0 g	4.0 g	6.0 g
175 mm	4.0 g	7.0 g	10.0 g
200 mm	5.0 g	10.0 g	15.0 g
250 mm	11.0 g	23.0 g	35.0 g
300 mm	16.0 g	32.0 g	48.0 g

INCORPORATION: Plant Nutrient Requirements / Uniformly mix PLANTING BED / FIELD: Plant Nutrient Requirements (incorporate) nursery fertiliser into potting media as follows:

Kilograms per Cubic Metres			
LOW 2kg	MED 3.5kg	HIGH 5kg	

(incorporate if possible or use lower rates) as follows:

Kilograms per 100 SQ. Metres				
LOW 6kg	MED 12kg	HIGH 18kg		



KEEP OUT OF REACH OF CHILDREN WARNING **CAUSES SKIN IRRITATION.** CAUSES EYE IRRITATION. MAY CAUSE RESPIRATORY IRRITATION.

PRECAUTIONARY STATEMENTS: Avoid breathing dust. Wash thoroughly after handling. Use only outdoors or in a wellventilated area. Call a poison poison information centre (Phone 131126) / doctor if you feel unwell. Specific treatment see First Aid section on this label.

PERSONAL PROTECTIVE EQUIPMENT: Wear protective gloves / protective clothing / eye protection / face protection. Take off contaminated clothing and wash before reuse.

FIRST AID: IF ON SKIN Wash with plenty of water. If skin irritation occurs, get medical advice / attention, IF INHALED Remove person to fresh air and keep comfortable for breathing, IF IN EYES Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical advice / attention.

STORAGE AND DISPOSAL - STORE in a well-ventilated place. Keep container tightly closed. Store locked up. DISPOSAL Dispose of contents / container in accordance with local / regional / national regulations.

APPLICATION PRECAUTIONS:

- Trial before use of this product under your local growing conditions, application methods, and desired rates. Avoid application to plants under stress.
- If mixed media is not used within one week, leach thoroughly before using.
- · Product left in media for more than one week will lose longevity resulting in reduced release time and wasted controlled release fertiliser.
- Avoid the use of media processing equipment that could change the integrity of the polymer coating.
- · Avoid mounding of fertiliser against base of plant.
- Iron and other plant nutrients can cause staining of cement.
- Keep away from pools, ponds, and other bodies of water.
- When using potting media with higher cation exchange capacities use lower recommended rates of this formulation.
- When using supplemental liquid feed reduce the rate of this formulation accordingly.
- Do not incorporate into media prior to steam sterilization.
- This product is not recommended for dibble applications.
- . To avoid buildup of soluble salts, occasional leaching may be necessary.
- . CAUTION: Application of fertiliser materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) that are toxic to ruminant

CONDITIONS OF SALE: ISP shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on ISP's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of ISP has any authority to add to or alter these conditions.





^{**18.00%} slowly available urea nitrogen from polymer-coated urea.