

Turf

Apply in sufficient volume of water to achieve through coverage of all foliage. In turf this can generally be achieved with spray volumes of 4-8 L/100m². Ensure even application in order to achieve even disease control. Avoid overlapping sprays.

TRADE ADVICE:
Export of treated produce

Grapes

While Maximum Residue Limits (MRLs) have been set in many major wine export destinations, some export destinations have not finalised MRL applications. For further information regarding export tolerances please contact your winery, INDIGO Specialty Products Pty Ltd representative or the Australian Wine Research Institute.

Other Crops

While Maximum Residue Limits (MRLs) have been set in many major export destinations, it should be noted that MRLs or import tolerances may not be established in all export destinations. For further information regarding export tolerances please contact your export organisation or INDIGO Specialty Products Pty Ltd representative.

Fungicide Resistance Warning

GROUP 11 FUNGICIDE

AZO-FORCE 250 SC Fungicide is a member of the Strobilurin and related products group of fungicides. For fungicide resistance management the product is a Group 11 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, INDIGO Specialty Products Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

Resistance Management in Turf

AZO-FORCE 250 SC Fungicide is best used in a preventative disease control program. Reliance on curative use of AZO-FORCE 250SC Fungicide could lead to resistance. **As a precaution, DO NOT apply AZOXYSTROBIN Fungicides for more than 1/3 of the total fungicide applications in one year.** If consecutive applications of AZO-FORCE 250 SC Fungicide are used they must be followed by at least the same number of applications of fungicides from a different group before AZO-FORCE 250 SC Fungicide is used again that year.

PRECAUTION

Re-Entry Period - Turf

DO NOT enter treated areas until the spray has dried unless wearing chemical resistant gloves and footwear.

PROTECTION OF LIVESTOCK

DO NOT graze treated turf/lawn or feed turf/lawn clippings from any treated area to poultry or livestock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

HIGHLY TOXIC TO AQUATIC LIFE. **DO NOT** contaminate dams, waterways or drains with the chemical or used containers.

DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift on adjacent areas, particularly wetlands, waterbodies or watercourses.

DO NOT apply when heavy rains or storms are expected within 4 hours.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty container or product.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes.

When opening the container and preparing spray wear:

- cotton overalls, buttoned to the neck and wrist (or equivalent clothing).
- elbow-length PVC gloves.

IF APPLYING BY HAND WEAR:

- cotton overalls, buttoned to the neck and wrist over normal clothing
- elbow-length PVC gloves.

Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

FIRST AID: IF POISONING OCCURS, CONTACT A DOCTOR OR POISONS INFORMATION CENTRE. PHONE: AUSTRALIA 131126, NEW ZEALAND 0800 764 766.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

NOTICE TO BUYER

INDIGO Specialty Products Pty Ltd makes no warranty expressed or implied, concerning the use of this product other than that indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

CROP	DISEASE	RATE/100L	WHP	CRITICAL COMMENTS
Poppies	Downy Mildew	750 ml/ha	6 weeks	Apply AZO-FORCE 250 SC Fungicide in a preventative fungicide program containing fungicides from a different chemical group. Apply AZO-FORCE 250 SC Fungicide preventatively before disease symptoms appear. Ensure thorough spray coverage. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to AZO-FORCE 250 SC Fungicide develop. DO NOT use AZO-FORCE 250 SC Fungicide curatively. Apply a maximum of 2 applications of AZO-FORCE 250 SC Fungicide per crop.
Potatoes	Early blight (Target Spot) (<i>Alternaria solani</i>)	300 to 400 ml/ha		Apply the higher rates when climatic conditions favour Early Blight or Late Blight infection and in crops with large canopies. Application AZO-FORCE 250 SC Fungicide may be applied by ground or aerial application equipment in potatoes. Aerial application may be used only for Early Blight (Target Spot) control. Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the potatoes. Ground Application: A volume of 200 to 300 L/ha is suggested at the start of the season, increasing to 500 to 600 L/ha in a vigorous crop at full canopy. Aerial Application (Early Blight only): A volume of 30 to 40 L/ha is recommended. Spray Interval Consecutive applications should be applied at 7 to 14 day intervals at any time between early shoot growth and 14 days before harvest. Use the recommended shorter application interval in the following circumstances: 1. Under humid weather conditions which are favourable for Early or Late Blight infection. 2. When there is rapid vegetative growth during the early part of the crop cycle. 3. At the first sign of Late Blight infection. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to AZO-FORCE 250 SC Fungicide develop. Apply AZO-FORCE 250 SC Fungicide in a protective spray program. DO NOT wait until disease levels have built up to make applications as this reduces the effectiveness of control and increases the risk of resistance development. As a precaution, DO NOT apply more than 1/3 of the total fungicide sprays per crop as AZO-FORCE 250 SC Fungicide. A maximum of 3 consecutive applications of AZO-FORCE 250 SC Fungicide are to be applied. They must be preceded and followed by at least 3 applications of fungicide(s) from a different fungicide group(s) such as Chlorothalonil, before AZO-FORCE 250 SC Fungicide is used again in that crop. Where Late Blight infection has occurred it is recommended that single sprays of AZO-FORCE 250 SC Fungicide be alternated with 2 sprays of Chlorothalonil or a fungicide(s) from another group(s). Where crops are grown successively alternation should continue between crops.
	Late Blight (<i>Phytophthora infestans</i>)	500 to 600 ml/ha		
	Soil borne: Black Scurf (<i>Rhizoctonia solani</i>) Silver Scurf* (<i>Helminthosporium solani</i>) *Suppression only.	5 to 10ml/ 100m of row		Apply once as an in-furrow spray at planting. Mount the spray nozzle so the spray is directed into the furrow as a 15 to 20 cm band just before the seed is covered. Use the higher rate of AZO-FORCE 250 SC Fungicide where higher levels of disease occur. Use the lower rate where lower levels of disease occur or where less disease control is required. Apply in 1 to 3 L of water/100m of row. Ensure the water volume used is not so high as to wash off any seed treatments previously applied to seed. DO NOT apply AZO-FORCE 250 SC Fungicide if conditions or seed quality favour bacterial rots as these diseases may be aggravated if seed comes into contact with additional moisture. DO NOT apply AZO-FORCE 250 SC Fungicide if planting in hot, sandy soils as bacterial rots may be aggravated. DO NOT mix with other products when applying as an in-furrow spray.
Tomatoes Except greenhouse	Early Blight (Target Spot) (<i>Alternaria solani</i>)	400 mL/ha or 40mL/100 L	1 day	Application Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the tomatoes and the method of trellising which influences canopy volume. In the case of dilute spraying (mL/100 L) apply in the range of 400 to 500 L/ha after transplanting and increase to 800 to 1000 L/ha at full canopy. In the case of fully trellised tomatoes at full canopy, application volumes should be increased to 1500 L/ha to achieve these results with high volume spraying. For Late Blight and Sclerotinia control use the higher rates when climatic conditions are humid and mild, which favours disease infection. Spray Interval Consecutive applications should be applied at 7 to 14 day intervals commencing soon after transplanting and continuing up to fruit maturity. Use the recommended shorter application interval in the following circumstances: 1. Under humid weather conditions which are favourable for disease infection. 2. When there is rapid vegetative growth during the early part of the crop cycle. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to AZO-FORCE 250 SC Fungicide develop. Apply AZO-FORCE 250 SC Fungicide in a protective spray program. DO NOT wait until disease levels have built up to make applications as this reduces the effectiveness of control and increases the risk of resistance development. As a precaution, DO NOT apply more than 1/3 of the total fungicide sprays per crop as AZO-FORCE 250 SC Fungicide. A maximum of 3 consecutive applications of AZO-FORCE 250 SC Fungicide are to be applied. They must be preceded and followed by at least 3 applications of fungicide(s) from a different fungicide group(s) such as Chlorothalonil, before AZO-FORCE 250 SC Fungicide is used again in that crop. Where Late Blight infection has occurred it is recommended that single sprays of AZO-FORCE 250 SC Fungicide be alternated with 2 sprays of Chlorothalonil or a fungicide(s) from another group(s). Where crops are grown successively alternation should continue between crops.
	Late Blight (<i>Phytophthora infestans</i>), Sclerotinia (<i>Sclerotinia minor</i>)	500 to 600 mL/ha or 50 to 60 mL/100 L		
NON-CROP SITUATIONS				
SITUATION	DISEASE	RATE	CRITICAL COMMENTS	
Turf			DO NOT use in the home garden DO NOT apply by aerial application methods DO NOT use this product through any type of irrigation system or ultra-low volume spray system	
	Anthracnose (<i>Colletotrichum graminicola</i>) Grey leaf spot (<i>Pyricularia grisea</i>) Helminthosporium disease (<i>Bipolaris</i> spp., <i>Drechslera</i> spp., <i>Exserohilum</i> spp.) Red Thread (<i>Laerisaria fruciformis</i>) Winter fusarium (<i>Microdochium Patch</i>) (<i>Microdochium nivale</i> / <i>Fusarium nivale</i>) <u>Brown patch (<i>Rhizoctonia solani</i>)</u>	12-23 mL /100m² applied in 4-8 L (2.3 L /ha)	Apply in sufficient volume of water to achieve thorough coverage of all foliage. Apply AZO-FORCE 250 SC Fungicide in a preventative fungicide program containing fungicides from different chemical groups (such as propiconazole and chlorothalonil). Begin applications before symptoms occur when conditions favour disease and continue applications while conditions remain favourable for disease development.Apply 2 consecutive applications at 14 to 21 day intervals. Use the shorter spray interval when climaticconditions remain favourable for disease development over prolonged periods. Spray interval Apply no more than 2 consecutive AZO-FORCE 250 SC Fungicide applications at 14-21 days intervals for Helminthosporium and at 14-28 day intervals for the other diseases. Use the shorter spray interval or apply the higher rate when climatic conditions remain favourable for disease development over prolonged periods. Use the shorter spray interval and apply the higher rate when conditions for disease development are optimum. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to AZO-FORCE 250 SC Fungicide develop. AZO-FORCE 250 SC Fungicide is best used in a preventative disease control program. Reliance on curative use of AZO-FORCE 250SC Fungicide could lead to resistance. As a precaution, DO NOT apply Fungicides for more than 1/3 of the total fungicide applications in one season. If consecutive applications of AZO-FORCE 250 SC Fungicide are used they must be followed by at least the same number of applications of fungicides from a different group before AZO-FORCE 250 SC Fungicide is used again in the current or following season.	
	Pythium Leaf Blight, Pythium Root Rot, Seedling Damping Off (<i>Pythium spp.</i>)	12-23 mL /100m² applied in 4-8 L (2.3 L /ha)	Apply in sufficient volume of water to achieve thorough coverage of all foliage (Pythium leaf blight). Wash in within 1 hour after application (Pythium root rot and Seedling damping off). Apply AZO-FORCE 250 SC Fungicide in a preventative fungicide program containing fungicides from different chemical groups Begin applications when conditions first favour disease development and before disease is present. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to AZO-FORCE 250 SC Fungicide develop. AZO-FORCE 250 SC Fungicide is best used in a preventative disease control program. Reliance on curative use of AZO-FORCE 250SC Fungicide could lead to resistance. As a precaution, DO NOT apply AZOXYSTROBIN Fungicides for more than 1/3 of the total fungicide applications in one season. If consecutive applications of AZO-FORCE are used they must be followed by at least the same number of applications of fungicides from a different group before AZO-FORCE is used again in the current or following season.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION