

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Contra M Duo

HERBICIDE

ACTIVE CONSTITUENTS:

340 g/L MCPA present as the dimethylamine salt

80 g/L DICAMBA present as the dimethylamine salt

GROUP	I	HERBICIDE
-------	---	-----------

For the Control of Certain Broadleaf Weeds in Winter Cereals, Pastures, Turf and Non-Crop Areas as per the Directions for Use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT

indigo
SPECIALTY PRODUCTS

Indigo Specialty Products Pty Ltd
ABN 15 631 459 660
3/49 Donaldson Road
Rocklea QLD 4106
Ph: 0491 201 694

Contents
20 L



Contra M Duo

HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in accordance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour. Repeated exposure may cause allergic disorders. When opening container and preparing spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves, face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor, or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce CONTRA M DUO Herbicide is available from Indigo Specialty Products Pty Ltd on request.

CONDITIONS OF SALE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

	IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE
	SPECIALIST ADVICE IN AN EMERGENCY ONLY 0417 653 654 ALL HOURS – AUSTRALIA WIDE



20L

Batch No.

Date of
Manufacture

APVMA Approval No.: 88327/122176

Contra M Duo

HERBICIDE

DIRECTIONS FOR USE

Restraints
DO NOT spray when rain seems likely to occur with 4 hours.
DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging.
DO NOT spray outside recommended crop growth as crop damage may result.
DO NOT apply to crops or pastures with clover, lucerne or medics.
DO NOT apply to legume pastures.
DO NOT apply to plants suffering from stress.

Crop & Situation	Weeds	State	Rate/ha	Critical Comments
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Zadoks 21)	Capeweed, Charlock, Common Ice-Plant, seedling Docks, Doublegee (Three cornered Jack, Spiny Emex) Hexham Scent (Melilotus), Mustards, Solider Thistle, Sorrel Seedlings, Tares, Tree Hogweed, Turnip Weed (<i>Rapistrum rugosum</i>), Vetches, Ward's Weed, Wild Radish, Wild Turnip (<i>Brassica tournefortii</i>), Wireweed (Hogweed)	All States	1 L	Spray when most weeds are in the 2 to 4 leaf stage. In some instances suppression only of Wireweed (Hogweed), Capeweed and Wild Radish will occur.
	Seedling Cotton Thistle, Fat-Hen	Tas		
	Cotton Thistle to bud formation	Tas	1.7 L	Spray when most weeds are still in the rosette or seedling stage.
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)	Mintweed, New Zealand Spinach	Qld, NSW		
	Capeweed, Charlock, Chickweed, Climbing Buckwheat (Black Bindweed), Clover, Common Ice-Plant, Seedling Docks, Doublegee	Qld, NSW, Tas, WA, Vic	1.4 L	N.B. Wild radish may need to be resprayed (with 2,4-D or MCPA) if future germinations occur after spraying. In some instances, suppression only of wireweed (hogweed), hoary cress and climbing buckwheat will be obtained.
	(Three cornered Jack, Spiny Emex), Fat-Hen, Hexham Scent (Melilotus), Seedling Hoary Cress, Mouse-eared Chickweed, Mustards, Narrow-leaved Plantain, Saffron Thistle, Spurry, Swinecress, Tares, Tree Hogweed, Turnip Weed (<i>Rapistrum rugosum</i>), Variegated Thistle, Vetch, Wild Radish, Wild Turnip (<i>Brassica tournefortii</i>), Wireweed (Hogweed)	SA		
	Volunteer Peas	SA	1 L plus 60 mL/ha dicamba (500 g/L)	Apply when most peas are 5-7cm high.
Oats, Wheat From early tillering stage (Zadoks 21) until the fully tillered stage before jointing occurs (Zadoks 30)			1 L/ha plus 120 mL/ha dicamba (500 g/L)	

SDS
Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce CONTRA M DUO Herbicide is available from Indigo Specialty Products Pty Ltd on request.

CONDITIONS OF SALE:
Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

FIRST AID
If poisoning occurs, contact a doctor, or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DIRECTIONS
Harmful if absorbed by skin contact, inhaled or swallowed. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point.
Store in the closed, original container in a dry, cool, well-ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Minimise spray drift by using low pressure and nozzles which DO NOT give a fine droplet size. DO NOT use in high wind.
DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into use. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops.
DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures.
Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, vines, lupins, fruit trees and ornamentals. empty containers or product.

GENERAL INSTRUCTIONS
MIXING
Add the required amount of CONTRA M DUO directly to the tank and mix well.

APPLICATION
For Boom Application: Apply by a properly calibrated boom spray using not less than 50 litres of water per hectare unless indicated otherwise in Critical Comments column.
For Aerial Application: Apply at least 25 litres total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.
For High Volume Application: Apply minimum 1500 litres per hectare.
For Knapsack Application: A 15 litre knapsack treats 150 square metres.
NOTE: Spray when weeds are young and actively growing. A temporary wilting may be evident in crops after application. If seasonal conditions are unusually late or dry seek further advice as damage may result from spraying under these conditions. Growers should seek advice before spraying recently released cereal varieties. There is nil tolerance of pea seed contaminants in cereal grain and less than 100% control of Volunteer Peas could result in dockage.

Situation	Weeds Controlled	State	Rate			Critical Comments
			Per ha	Per 100 L	Per 15 L Knap-sack	
Grass Pastures	Bathurst Burr, Caltrop (Yellow Vine), Chicory, Crofton Weed, Cudweed, Dandelion, Deadnettle, Fleabanes, Fumitory, Groundsel, Heliotrope, Inkweed, Mintweed, Nettles, Noogoora Burr, Pennycress, Potato Weed, Ragwort, Ripplewort, Shepherd's Purse, Smartweed, Soft Rush, Stinkwort, Thistles including Cockspur, Scotch (Spear), Slender, Star, Variegated, Wireweed (Hogweed)	All States	2.8-4 L	190 mL – 270 mL	60 mL	Use the higher rate on larger weeds.
	Cotton Thistle	Tas				
	Buttercups, Hoary Cress seedlings, Horehound, Khaki Weed, Purple-top, Veined Verbena	All States	4-7 L	470 mL	100 mL	When necessary a second treatment may be made after 4 weeks.
Turf	Bindy-Eye, Capeweed, Catsear, Chickweed, Clovers, Cotula, Creeping Oxalis, Cudweeds, Dandelion, Dock, Fleabanes, Fumitory, Jo-Jo (Onehunga), Lamb's Tongue, Medics, Mustards, Narrow-leaved Plantain, Pearlwort, Pennywort, Peppergrass, Radish, Scarlet Pimpernel, Slender Thistle, Soldier Thistle, Spear Thistle, Toad Rush, Variegated Thistle, Wild Turnip, Wireweed (Hogweed)	All States	Golf Tee, Golf Green: 100 mL in 15 L water per 150 m² Bowling Green: 900 mL in 200 L water per 1400 m² Large Turf Area: (Field, Park, Golf Fairway) 6.5 L in 250-400 L water per 10,000 m² (1 hectare)			Apply to actively growing weeds and moist soil. DO NOT use on Lippia or Strawberry Clover or Buffalo grass. DO NOT mow turf for 2 days before or after application or fertilize within two weeks of spraying.
	Fat-Hen, Purple – Top, Red Flowered Mallow, Stagger Weed, Swinecress	Qld, NSW, Tas, SA, WA,				
	Artichoke, Bindweed, Fennel, Hemlock, Thistle seedlings	All States	5.2 L	350 mL	80 mL	
Non-Crop Situations						

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION
HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED

PLANTS
Spray equipment must be calibrated accurately prior to working in crops.
Minimise spray drift by using low pressure and nozzles which DO NOT give a fine droplet size. DO NOT use in high wind.
DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into use. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops.
DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures.
Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, vines, lupins, fruit trees and ornamentals. empty containers or product.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET
COMPATABILITY
CONTRA M DUO is compatible with flowable terbutryn.

RESISTANT WEEDS WARNING
ProForce CONTRA M DUO Herbicide is a member of the Benzoic acid and Phenoxy groups of herbicides. CONTRA M DUO has the disruptors of plant cell growth mode of action. For weed resistance management CONTRA M DUO is a Group I Herbicide. Some naturally occurring weed biotypes resistant to CONTRA M DUO and other Disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CONTRA M DUO or other Group I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Indigo Specialty Products Pty Ltd accepts no liability for any losses that may result from the failure of CONTRA M DUO to control resistant weeds.



Indigo Specialty Products ABN 15 631 459 660
3/49 Donaldson Road, Rocklea QLD 4106

Ph: 0491 201 694

APVMA Approval No.: 88327/122176

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Contra M Duo

HERBICIDE

ACTIVE CONSTITUENT: 340 g/L MCPA present as the dimethylamine salt
80 g/L DICAMBA present as the dimethylamine salt

GROUP I

HERBICIDE

For the Control of Certain Broadleaf Weeds in Winter Cereals, Pastures, Turf and Non-Crop Areas as per the Directions for Use.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.



Indigo Specialty Products Pty Ltd
ABN 15 631 459 660
3/49 Donaldson Road, Rocklea QLD 4106