## **DANGEROUS POISON**

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## **HERBICIDE**

**ACTIVE CONSTITUENTS:** 

200 g/L BROMOXYNIL present as the n-octanoyl ester 200 g/L MCPA present as the ethyl hexyl ester

**SOLVENT:** 

343 g/L LIQUID HYDROCARBON

GROUP 6 4 HERBICIDE

For the control of weeds in wheat, oats, barley, cereal rye, triticale, linseed, grass pastures and turf as specified in the Directions For Use table

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT



# Weed Blast MA

### **HERBICIDE**

### STORAGE AND DISPOSAL

This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

### Refillable containers

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### **SAFETY DIRECTIONS**

Product is poisonous if inhaled or swallowed. Attacks the eyes and will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting.

### SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce WEED BLAST MA Herbicide is available from Indigo Specialty Products Pty Ltd on request.

### **CONDITIONS OF SALE**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.



**0417 653 654**ALL HOURS – AUSTRALIA WIDE



Batch No.

Date of Manufacture

APVMA Approval No.: 70206/RV2024

# leed Blast MA

### **DIRECTIONS FOR USE** RESTRAINTS

DO NOT spray if weeds are wilted by dry or cold weather or if crops are at a susceptible stage, or if rain is likely within 4 hours.

DO NOT apply if frosts are imminent.

DO NOT apply to diseased or frost affected crops or turf.

### SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

FLAT SIZE: 330 MM WIDE X 446 MM DEEP

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

SITUATION & CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Cereal rye, Oats and Triticale; Linseed; Grass Pastures	Amsinckia (Amsinckia spp.), Capeweed (Arctotheca calendula), Chamomile (Matricaria matricarioldes), Charlock (Sinapis arvensis), Corn gromwell (Ironweed, Sheepweed) (Buglossoides arvensis), Fat hen (Chenopodium album), Field madder (Sherardia arvensis), Hexham scent (King Island Melilot) (Melilotus indica), Horned poppy (Glaucium flavum), Indian hedge mustard (Sisymbrium orientale), Lesser swinecress (Coronopus didymus), Mexican poppy (Argemone ochroleuca), Mintweed (Salvia reflexa), Paterson's curse (Salvation Jane) (Echium plantagineum), Rough poppy (Papaver hybridum), Saffron thistle (Carthamus lanatus), Shepherd's purse (Capsella bursapastoris), Slender thistles (Carduus tenuiflorus, C.pycnocephalus), Tree hogweed (Polygonum patulum), Turnip weed (Giant Mustard) (Rapistrum rugosum), Wild radish (Raphanus raphanistrum), Wild turnip (Brassica tournefortin)	Plants up to the 6 leaf stage but not more than 50 mm in diameter Plants up to the 8 leaf stage but not more than 75 mm in diameter.	WA only QId, NSW, Vic, Tas, SA, NT, ACT only WA only	750 mL/ha 1.4 L/ha 2.0 L/ha 1.4 L/ha	CROP STAGE: Wheat, Barley, Cereal Rye, Oats and Triticale: 3 leaf to fully tillered stage (Z13-30) for rates of 750 m.t. to 1.4 L/ha; 5 leaf to fully tillered stage (Z15-30) for rate of 2 L/ha (NSW, ACT, Vic, SA, WA, Tas only); or early tillering to fully tillered stage (Z21-30) (Qid only).  Linseed: 50 to 150 mm high. D0 NOT apply to linseed by aircraft.  Yell f spraying Black bindweed when wheat, barley, cereal rye, oats, triticale or linseed is more than 300 mm high, increased spray volume to 200 L/ha.  For optimal control, apply only when the weeds are actively growing.  In the early stages following spraying, temporary wilting of crop and pasture may occur.  Crop scorch may occur, particularly on oats if treatment is delayed to later stages of growth. Refer to the General Instructions.  Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days
	Black bindweed <sup>¥</sup> (Climbing buckwheat) ( <i>Fallopia convolvulus</i> ), Common peppercress ( <i>Lepidium africanum</i> ), Mustards ( <i>Sisymbrium</i> spp.), Purple Calandrinia ( <i>Calandrinia menziesii</i> )	Plants up to the 4 leaf stage but not more than 35 mm in diameter.	WA only	1 L/ha	after application. DO NOT apply to crops or pastures undersown with lucerne, clover or medics unless some damage is acceptable.
		Plants up to the 6 leaf stage but not more than 50 mm in diameter	All States	1.4 L/ha	
		Plants up to the 8 leaf stage but not more than 75 mm in diameter.		2 L/ha	

available from Indigo Specialty Products Pty Ltd on request. A safety data sheet for ProForce WEED BLAST AM Herbicide is Additional information is listed in the safety data sheet (SDS). SDS

swallowed, do NOT induce vomiting. Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. It If poisoning occurs contact a doctor or Poisons Information **GIA T2811** 

ciorning.

**ANGEROUS POISON** 

SAFETY DIRECTIONS

day's use, wash gloves, face shield or goggles and contaminated hands, arms and face thoroughly with soap and water. After each water. After use and before eating, drinking of smoking, wasn shield or goggles. If product in eyes, wash it out immediately with wrist, a washable hat and elbow-length PVC gloves and face brepared spray wear cotton overalls buttoned to the neck and NOT inhale spray mist. When preparing the spray and using the and will irritate the skin. Avoid contact with eyes and skin. DO Product is poisonous if inhaled or swallowed. Attacks the eyes

**CONDITIONS OF SALE** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.



Indigo Specialty Products ABN 15 631 459 660 3/49 Donaldson Road, Rocklea QLD 4106 Ph: 0491 201 694

APVMA Approval No.: 70206/RV2024A

Envirodrum remains the property of Indigo Specialty Products Pty return the empty Envirodrum to the point of purchase. The When the contents of the Envirodrum have been used, please rinsed with clean water and drained accordingly.

Matic coupler, delivery system and hoses are disconnected, triple matter. After each use of the product, please ensure that the Micro NOT contaminate the Envirodrum with water or any other foreign NOT tamper with the Micro Matic valve or the security seal. DO area. DO NOT store for prolonged periods in direct sunlight. DO опуна зеаеа Епуноагит та соог, мен-чепиг Envirodrum 110 L Mini Bulk Returnable Container

return to point of supply for refill or storage.

contents fully into application equipment. Close all valves and srea. DO NOT store for prolonged periods in direct sunlight. Empty Store in the closed, original container in a cool, well-ventilated Refillable containers



THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. PRODUCT. **LIQUID HYDROCARBON** GROUP 6 4. HERBICIDE

Rocklea QLD 4106

e control of weeds necified in the Directi

SITUATION & CROP WEEDS CONTROLLED **WEED STAGE** STATE RATE CRITICAL COMMENTS CROP STAGE: Wheat, Barley Fumitory (Fumaria spp.), Three-cornered Qld, NSW, 1.4 L/ha Plants up to the 3 Cereal rye, Oats and Jack (Doublegee, Spiny emex) (Emex leaf stage but not ACT, Vic. Wheat, Barley, Cereal rye, Oats and Triticale: 3 leaf to fully tillered stage (Z13-30) for rates of 750 mL to 1.4 L/ha; 5 leaf to fully tillered stage (Z15-30) for rate Triticale: Linseed: more than 50 mm Tas, SA only **Grass Pastures** in diameter of 2.0 L/ha (NSW, ACT, Vic, SA, WA, Tas only); or early 750 mL/ha Plants up to the 4 WA only tillering to fully tillered stage (Z21-30) (Qld only). leaf stage but not Linseed: 50 to 150 mm high. DO NOT apply to linseed more than 60 mn in diameter Qld, NSW, Plants up to the 5 2.0 L/ha For optimal control, apply only when the weeds are Vic, Tas, SA, NT leaf stage but not actively growing. more than 75 mm In the early stages following spraying, temporary ACT only wilting of crop and pasture may occur. Plants up to the 6 WA only 1.4 L/ha Crop scorch may occur, particularly on oats if leaf stage but not treatment is delayed to later stages of growth. Refer to more than 85 mm the General Instructions in diameter Variegated thistle (Silybum marianum) Plants up to the  $4\,$ 750 mL/ha Wireweed (Hoaweed) leaf stage but not Qld, NSW 1.4 L/ha (Polygonum aviculare) more than 35 mm Vic, Tas, in diameter SA. NT. ACT Avoid application when maximum daily temperatures Plants up to the 6 only above 20°C occur, or are likely to occur for a few days leaf stage but not after application. DO NOT apply to crops or pastures more than 50 mm undersown with lucerne, clover or medics unless in diameter WA only 1.4 L/ha some damage is acceptable 750 mL/ha Mountain sorrel (Oxalis acetosella), Plants up to the 6 Three-horned bedstraw (Galium leaf stage Where Wireweed (Hogweed) occurs in red soils of low fertility, it has been found to be less susceptible. In Plants up to the 8 1.4 L/ha these circumstances use 2 L/ha. leaf stage Cleavers (Galium aparine) 2-4 stem stage 750 mL/ha and 1-3 whorls of leaves per stem Apply under good growing conditions. Avoid spraying Wheat, Barley Common sowthistle (Sonchus oleraceus) Cereal rye, Oats when soil is dry. DO NOT spray under frosty conditions Triticale In early stages following spraying, temporary wilting of crop may occur. Avoid application when maxim daily temperatures above 20°C or are more likely to occur for a few days after application. DO NOT apply to crops undersown with lucerne, clover or medic unless some damage is acceptable. Crop stages: Wheat, Barley, Oats, Triticale: - 3 leaf to Up to 4 leaves Northern 1.4 L NSW and full tillering (Z13-30). Use as early as possible to avoid Qld only shading of weeds by the crop. Strictly for use on small weeds early in crop before shading out. If Group 3 resistance is suspected, or weeds have grown through a previous application of chlorsulfuron apply 2.1 L/ha. Use a minimum spray volume of 70 L/ha 4 to 8 leaves 2.1 L Crop stages: Wheat, Barley, Oats and Triticale: -3 leaf to full tillering (Z13-30). Use as early as possible to avoid shading of weeds by the crop. Turf, established Bindy-eye, Jo-Jo (Onehunga), Cudweed All States Apply evenly over the area to be treated. DO NOT mow Couch (incl. Dandelion, Catsear, Plantains, Thistles ml /100 I turf for 2 days after treatment

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

wate

OR 60 mL/100

WITHHOLDING PERIOD: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION

empty containers or product.

Local, State or Territory government regulations. DO NOT burn desirable vegetation and tree roots, in compliance with relevant specifically marked and set up for this purpose clear of waterways, empty packaging 500 mm below the surface in a disposal pit approved waste management facility is not available, bury the backaging to an approved waste management facility. If an If not recycling, break, crush or puncture and deliver empty

Oxalis, Capeweed, Creeping buttercup

Fleabane, Clover (not Tas.)

clean containers to recycler or designated collection point. undiluted chemicals on site. If recycling, replace cap and return before disposal. Add rinsings to spray tank. DO NOT dispose of store for prolonged periods in direct sunlight. Triple rinse containers closed, original container in a cool, well-ventilated area. DU NUI children, animais, 1000, reedsturis, seed and refulisers. Store in This product must be stored in a locked room or place away from

JASO9SIQ QNA 3DAROTZ

If in doubt, contact your nearest Department of Agriculture. breeding stock. DO NOT graze horses or pigs on Paterson's curse. thistles predominate in the pasture. Avoid grazing with young or esbecially where capeweed, Paterson's curse and variegated from causes such as nitrate poisoning. Care should be taken ilitake of some weeds may result in stock poisoning and death Sprayed weeds may become more palatable to stock and a higher

Grazing Precaution PROTECTION OF LIVESTOCK

watercourses with the chemical or used containers. Dangerous to fish. DO NOT contaminate streams, rivers or PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

### **GENERAL INSTRUCTIONS**

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid Instructions.

### MIXING

Queensland blue).

Kikuyu, Paspalum,

Buffalo, Fescue. Bent & Rye lawns

To ensure even mixing, half fill spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of water. Agitate again before spraying

### **APPLICATION**

DO NOT use rain-drop or air-induction nozzles as coverage of weeds may not be adequate.

**Boom Sprayer:** Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom

above the crop to ensure that the spray is evenly distributed. Cereals: Apply 50-200 L of water per hectare. Use higher water volume (min. 100 L/ha) where the weed infestation is heavy, the crop cover is dense or when tank-mixing with DIMOP 500. Complete coverage of weeds is essential.

Linseed: Apply a minimum of 220 L of water per hectare. Turf: Apply a minimum of 500 L of water per hectare.

Aerial Application: Apply a minimum of 22 L of water per hectare or 30 L of water per hectare if tank-mixing with DIMOP 500. Markers should always move upwind before aircraft passes overhead. Application by micronaire equipment may give variable

and established plants may be damaged. or on days following treatment, may result in seedling mortality Note: Maximum daily temperatures at or above 20°C at application, by misting machines.

tomatoes, vines, lupins, fruit trees and ornamentals. DO NOT apply vapour movement onto susceptible crops such as cotton, tobacco, plants/crops, cropping lands or pastures. Avoid spray drift and equipment, that may cause spray to drift onto nearby susceptible DO NOT apply under weather conditions, or from spraying

### **STNAJ9** PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET

of WEED BLAST MA to control resistant weeds. accepts no liability for any losses that may result from the failure difficult to detect prior to use, Indigo Specially Products Pty Ltd Group 4 herbicides. Since the occurrence of resistant weeds is weeds will not be controlled by this product or other Group 6 and population if these herbicides are used repeatedly. These resistant The resistant individuals can eventually dominate the weed exist through normal genetic variability in any weed population. WEED BLAST MA and other Group 6 and Group 4 herbicides may onne naturally-occurring weed biotypes resistant to resistance management WEED BLAST MA is a Group 6 and Group Il sud disruptors of plant cell growth modes of action. For weed BLAST MA has both the inhibitors of photosynthesis at photosystem both the nitriles and the phenoxys groups of herbicides. WEED Herbicide is a member of

GROUP 6 4 HERBICIDE AM TSAJB GEEW 92707079 RESISTANT WEEDS WARNING

### **EQUIPMENT MAINTENANCE**

After use wash application equipment thoroughly with Spraymate Tank & Equipment Cleaner and rinse several times.

This product is physically compatible in the spray tank with one of the following herbicides or insecticides in wheat and barley only. METFURON, DIMOP 500, DICAM, MCPA, DIMETHOLINX, OMETHO-MITE, Endosulfan or CHLORPOS. When mixed with METFURON or DICAM, do not apply to cereals undersown with lucerne. When mixed with DIMOP 500, DO NOT exceed 1.4 L of this product per hectare. Observe all instructions on the mixture partner label. As formulations of other manufactures products are beyond the control of Indigo Specialty Products Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.